

# A minimum degree result for disjoint cycles and forests in bipartite graphs

DANHONG ZHANG    HONG WANG

*Department of Mathematics  
The University of Idaho  
Moscow  
Idaho 83844  
USA*

## Abstract

Let  $F = (U_1, U_2; W)$  be a forest with  $|U_1| = |U_2| = s$ , where  $s \geq 2$ , and let  $G = (V_1, V_2, E)$  be a bipartite graph with  $|V_1| = |V_2| = n \geq 2k + s$ , where  $k$  is a nonnegative integer. Suppose that the minimum degree of  $G$  is at least  $k + s$ . We show that if  $n > 2k + s$  then  $G$  contains the disjoint union of the forest  $F$  and  $k$  disjoint cycles. Moreover, if  $n = 2k + s$ , then  $G$  contains the disjoint union of the forest  $F$ ,  $k - 1$  disjoint cycles and a path of order 4.

## 1 Introduction

A set of graphs is called disjoint if no two of them have any vertex in common. Schuster [5] investigated the disjoint cycles and a forest in a graph. He proved the following result:

**Theorem A.** ([5], Theorem) *Let  $F$  be a forest on  $s$  edges without isolated vertices and let  $G$  be a graph of order at least  $3k + |V(F)|$  with minimum degree at least  $2k + s$ , where  $k$  and  $s$  are nonnegative integers. Then  $G$  contains the disjoint union of the forest  $F$  and  $k$  disjoint cycles.*

In this paper, we consider a similar problem in bipartite graphs. About the maximum number of disjoint cycles in a bipartite graph, H. Wang proved the following theorems:

**Theorem B.** ([7], Theorem 1) *Let  $G = (V_1, V_2; E)$  be a bipartite graph with  $|V_1| = |V_2| = n > 2k$ , where  $k$  is a positive integer. Suppose that the minimum degree of  $G$  is at least  $k + 1$ . Then  $G$  contains  $k$  disjoint cycles.*

**Theorem C.** ([7], Theorem 2) *Let  $G = (V_1, V_2; E)$  be a bipartite graph with  $|V_1| = |V_2| = n = 2k$ , where  $k$  is a positive integer. Suppose that the minimum degree*

of  $G$  is at least  $k + 1$ . Then  $G$  contains  $k - 1$  disjoint 4-cycles and a path of order 4 such that the path is disjoint from all the  $k - 1$  4-cycles.

This paper proves two theorems as follows:

**Theorem 1.** *Let  $F = (U_1, U_2; W)$  be a forest with  $|U_1| = |U_2| = s$ , where  $s \geq 2$ . Let  $G = (V_1, V_2, E)$  be a bipartite graph with  $|V_1| = |V_2| = n > 2k + s$ , where  $k$  is a nonnegative integer. Suppose that the minimum degree of  $G$  is at least  $k + s$ . Then  $G$  contains the disjoint union of the forest  $F$  and  $k$  disjoint cycles.*

**Theorem 2.** *Let  $F = (U_1, U_2; W)$  be a forest with  $|U_1| = |U_2| = s$ , where  $s \geq 2$ . Let  $G = (V_1, V_2; E)$  be a bipartite graph with  $|V_1| = |V_2| = n = 2k + s$ , where  $k$  is a nonnegative integer. Suppose that the minimum degree of  $G$  is at least  $k + s$ . Then  $G$  contains the disjoint union of the forest  $F$ ,  $k - 1$  disjoint cycles and a path of order 4.*

All graphs considered in this paper are finite simple graphs in standard terminology and notation from [1] except as indicated. Let  $G = (V, E)$  be a graph. For any  $u \in V$ , if  $G'$  is a subgraph of  $G$ , we define  $N(u, G')$  to be  $N_G(u) \cap V(G')$  and let  $d(u, G') = |N(u, G')|$ . If  $d(u, G) = 0$  or 1 we say that  $u$  is an isolated vertex or an endvertex of  $G$ , respectively. The minimum degree of  $G$  is denoted by  $\delta(G)$ . For a subset  $U$  of  $V$ ,  $G[U]$  is the subgraph of  $G$  induced by  $U$ . For two disjoint subgraphs  $G_1$  and  $G_2$  of  $G$ ,  $E(G_1, G_2)$  is the set of all edges of  $G$  between  $G_1$  and  $G_2$ . Let  $e(G_1, G_2) = |E(G_1, G_2)|$ , i.e.  $e(G_1, G_2) = \sum_{x \in V(G_1)} d(x, G_2)$ . A set of pairwise disjoint edges of  $G$  is called a matching in  $G$ . If  $M$  is a matching with the property that every vertex of  $G$  is incident with an edge of  $M$ , then  $M$  is called a perfect matching in  $G$ . The disjoint union of two graphs  $S$  and  $T$  is denoted by  $S \dot{\cup} T$ . We use the symbol  $\bigcirc^k$  to denote the disjoint union of  $k$  cycles; for  $k = 1$  we simply write  $\bigcirc$  instead of  $\bigcirc^1$ . An embedding of a graph  $H$  into a graph  $G$  is an injective mapping  $\sigma : V(H) \rightarrow V(G)$  so that for every edge  $xy \in E(H)$ , the edge  $\sigma(x)\sigma(y)$  is contained in  $E(G)$ . We write  $H \subseteq G$  or  $G \supseteq H$  if there is an embedding of  $H$  into  $G$ . For an embedding  $\sigma$  of  $H$  into  $G$  and a subgraph  $M$  of  $H$ , let  $\sigma(M)$  denote the image of  $M$  in  $G$ , i.e.,  $\sigma(M)$  is the subgraph of  $G$  with vertex set  $\{\sigma(x) : x \in V(M)\}$  and edge set  $\{\sigma(x)\sigma(y) : xy \in E(M)\}$ . We use  $(X, Y; E)$  to denote a bipartite graph with  $(X, Y)$  as its bipartition and  $E$  as its edge set. The length of a cycle  $C$  is denoted by  $l(C)$ , and a 4-cycle is a cycle of length 4. An acyclic graph is a graph without cycles.

## 2 Lemmas

For all lemmas listed below,  $G = (V_1, V_2; E)$  is a given bipartite graph.

**Lemma 2.1** ([7], Lemma 2.1) *Let  $C$  be a cycle of  $G$  and  $x$  a vertex of  $G$  not on  $C$ . Suppose  $d(x, C) \geq 2$ . Then either  $C$  is a 4-cycle or  $C + x$  contains a cycle  $C'$  such that  $l(C') < l(C)$ .*

**Lemma 2.2** ([7], Lemma 2.2) *Let  $C$  be a 4-cycle of  $G$ . Let  $x \in V_1$  and  $y \in V_2$  be two vertices not on  $C$ . Suppose  $d(x, C) + d(y, C) \geq 3$ . Then there exists  $z \in V(C)$  such that either  $C - z + x$  is a 4-cycle and  $yz \in E$ , or  $C - z + y$  is a 4-cycle and  $xz \in E$ .*

**Lemma 2.3** ([7], Lemma 2.3) *Let  $T$  be a tree of order at least 2 with a bipartition  $(X, Y)$  such that  $|Y| \geq |X|$ . Let  $p = |Y| - |X|$ . Then  $Y$  contains at least  $p + 1$  endvertices of  $T$ .*

**Lemma 2.4** ([7], Lemma 2.4) *Let  $P = x_1x_2x_3$  and  $Q = y_1y_2y_3$  be two disjoint paths of  $G$  with  $x_1 \in V_1$  and  $y_1 \in V_2$ . Let  $C$  be a 4-cycle of  $G$  such that  $C$  is disjoint from both  $P$  and  $Q$ . Suppose  $d(x_1, C) + d(x_3, C) + d(y_1, C) + d(y_3, C) \geq 5$ . Then  $G[V(C \cup P \cup Q)]$  contains a 4-cycle  $C'$  and a path  $P'$  of order 6 such that  $P'$  is disjoint from  $C'$ .*

**Lemma 2.5** ([7], Lemma 2.5) *Let  $C$  be a 4-cycle of  $G$ . Let  $uv$  and  $xy$  be two disjoint edges of  $G$  such that they are disjoint from  $C$ . Suppose  $d(u, C) + d(v, C) + d(x, C) + d(y, C) \geq 5$ . Then  $G[V(C) \cup \{u, v, x, y\}]$  contains a 4-cycle  $C'$  and a path  $P'$  of order 4 such that  $P'$  is disjoint from  $C'$ .*

**Lemma 2.6** ([7], Lemma 2.6) *Let  $C$  be a 4-cycle and  $P$  a path of order 4 in  $G$  such that  $P$  is disjoint from  $C$  and  $\sum_{x \in V(P)} d(x, C) \geq 6$ . Then either  $G[V(C \cup P)]$  contains two disjoint quadrilaterals, or  $P$  has an endvertex, say  $z$ , such that  $d(z, C) = 0$ .*

**Lemma 2.7** ([7], Lemma 2.7) *Let  $C$  be a 4-cycle and  $P$  a path of order  $s \geq 6$  in  $G$  such that  $C$  is disjoint from  $P$ . If  $\sum_{x \in V(P)} d(x, C) \geq s + 1$ , then  $G[V(C \cup P)]$  contains two disjoint cycles.*

**Lemma 2.8** ([7], Lemma 2.8) *Let  $s$  and  $t$  be two integers such that  $t \geq s \geq 2$  and  $t \geq 3$ . Let  $C_1$  and  $C_2$  be two disjoint cycles of  $G$  with lengths  $2s$  and  $2t$ , respectively. Suppose that  $\sum_{x \in V(C_2)} d(x, C_1) \geq 2t + 1$ . Then  $G[V(C_1 \cup C_2)]$  contains two disjoint cycles  $C'$  and  $C''$  such that  $l(C') + l(C'') < 2s + 2t$ .*

**Lemma 2.9** *Let  $F = (U_1, U_2; W)$  be a forest with  $|U_1| = |U_2| = s$ , where  $s \geq 1$ . Let  $G = (V_1, V_2; E)$  be a bipartite graph with  $|V_1| = |V_2| = n \geq s$  and  $\delta(G) \geq s$ . Then  $G \supseteq F$ .*

**Proof.** Without loss of generality, assume  $F$  is a tree. The lemma is trivial for  $s = 1$ . By Lemma 2.3, each of  $U_1$  and  $U_2$  contains an endvertex of  $F$ , say  $x$  and  $y$ , respectively. Let  $F' = F - \{x, y\}$ . By induction on  $s$ , there exists an embedding  $\sigma$  of  $F'$  in  $G$ . Suppose  $x_1x, y_1y \in W$  with  $\{x_1, y_1\} \subseteq V(F')$ . Since  $\delta(G) \geq s$ ,  $N(\sigma(x_1), G - V(\sigma(F'))) \neq \emptyset$  and  $N(\sigma(y_1), G - V(\sigma(F'))) \neq \emptyset$ , and it follows  $G \supseteq F$ .

**Lemma 2.10** *Let  $F = (U_1, U_2; W)$  be a forest in  $G$  with  $|U_1| = |U_2| = s$ , where  $s \geq 3$ . Let  $C = (A_1, A_2; B)$  be a cycle in  $G$  with  $|A_1| = |A_2| = t \geq 3$ , and  $C$  is disjoint from  $F$ . Suppose  $e(C, F) \geq 2ts - 4$ , then  $G[V(C \cup F)] \supseteq C' \dot{\cup} F$ , where  $C'$  is a 4-cycle.*

**Proof.** Since  $e(C, F) \geq 2ts - 4$ ,  $t \geq 3$  and  $s \geq 3$ , there exist  $\{x, y\} \subseteq V(C)$  with  $x \neq y$  and  $d(x, F) = d(y, F) = s$ . We may choose  $x$  and  $y$  such that  $x \in A_1$  and  $y \in A_2$ . Suppose this is not the case, say, for any  $z \in A_2$ ,  $d(z, F) \leq s - 1$ . Let  $C = z_1 z_2 \dots z_{2t} z_1$  with  $z_1 \in A_1$ . As  $e(C, F) \geq 2ts - 4$ , either  $d(z_1, F) = s$  or  $d(z_5, F) = s$ . If  $w \in N(z_2, F) \cap N(z_4, F)$ , then  $G[V(C \cup F)] \supseteq C' \cup F$ , where  $C'$  is the 4-cycle  $wz_2z_3z_4w$  and  $F \subseteq F - w + z_i$  for some  $i \in \{1, 5\}$  with  $d(z_i, F) = s$ . So we may assume  $N(z_2, F) \cap N(z_4, F) = \emptyset$ . Therefore  $d(z_2, F) + d(z_4, F) \leq s$ . Then  $e(C, F) \leq t(s - 1) + s(t - 1) = 2ts - t - s < 2ts - 4$ , a contradiction, hence the claim is true. Then we see that for any  $i \in \{1, \dots, t - 1\}$  with  $z_{2i+1} \neq x$ ,  $N(z_{2i}, F) \cap N(z_{2i+2}, F) = \emptyset$ , and  $N(z_2, F) \cap N(z_{2t}, F) = \emptyset$  if  $x \neq z_1$ , for otherwise  $G[V(C \cup F)] \supseteq C' \cup F$ , where  $C'$  is a 4-cycle. When  $t$  is even, it's easy to deduce that  $\sum_{i=1}^t d(z_{2i}, F) \leq s(t/2)$  and  $\sum_{i=1}^t d(z_{2i-1}, F) \leq s(t/2)$ . So  $2ts - 4 \leq e(C, F) \leq ts$ , implying  $st \leq 4$ , a contradiction. Similarly, when  $t$  is odd, we obtain  $e(C, F) \leq 2((t - 1)s/2 + s) < 2ts - 4$ , a contradiction.

**Lemma 2.11** *Let  $F = (U_1, U_2; W)$  be a forest in  $G = (V_1, V_2; E)$  with  $|U_1| = |U_2| = s$ , where  $s \geq 3$ . Let  $w$  and  $xy$  be two disjoint edges of  $G$  such that they are disjoint from  $F$ . Suppose  $d(u, F) + d(v, F) + d(x, F) + d(y, F) \geq 4s - 3$  and  $G[V(F)] = K_{s,s}$ . Then  $G[V(F) \cup \{u, v, x, y\}] \supseteq F \dot{\cup} P$ , where  $P$  is a path of order 4.*

**Proof.** As  $\sum_{t \in T} d(t, F) \geq 4s - 3$  where  $T = \{u, v, x, y\}$ , either  $N(u, F) \cap N(x, F) \neq \emptyset$  or  $N(v, F) \cap N(y, F) \neq \emptyset$ . Say the former holds, and let  $w \in N(u, F) \cap N(x, F)$ . For the same reason, either  $d(v, F) > 0$  or  $d(y, F) > 0$ . Say  $d(y, F) > 0$ . Clearly,  $G[V(F)] - w + y$  contains  $F$  since  $G[V(F)] = K_{s,s}$ . As  $vuw$  is a path of  $G$ , the lemma follows.

**Lemma 2.12** *Let  $F = (U_1, U_2; W)$  be a forest in  $G$  with  $|U_1| = |U_2| = s$ , where  $s \geq 3$ . Let  $P = x_1 x_2 \dots x_{2t}$  be a path in  $G$ , where  $t \geq 3$ . Suppose  $P$  is disjoint from  $F$ ,  $G[V(F)] = K_{s,s}$  and  $e(P, F) \geq 2t(s - 1) + 1$ . Then  $G[V(F \cup P)] \supseteq F \dot{\cup} \emptyset$ .*

**Proof.** Without loss of generality, suppose  $U_1 \subseteq V_1$ . Suppose that there exists  $v \in U_1$  such that  $v \in N(x_i, F) \cap N(x_{i+2}, F)$  for some  $i \in \{1, \dots, 2t - 2\}$ . Then  $vx_i x_{i+1} x_{i+2} v$  is a 4-cycle in  $G$ . If  $d(x_j, F) \geq 1$  for some  $x_j \in V(P) \cap V_1 - \{x_{i+1}\}$ , then  $G[V(F) - \{v\}] + x_j$  contains  $F$  and so the lemma holds. So we may assume  $d(x_j, F) = 0$  for all  $x_j \in V(P) \cap V_1 - \{x_{i+1}\}$ . It follows that  $2t(s - 1) + 1 \leq e(P, F) \leq ts + s$ , which implies  $(t - 1)(s - 2) - 1 \leq 0$ , a contradiction. So we may assume  $N(x_i, F) \cap N(x_{i+2}, F) = \emptyset$  and therefore  $d(x_i, F) + d(x_{i+2}, F) \leq s$  for all  $i \in \{1, \dots, 2t - 2\}$ . If  $t$  is odd, then  $2t(s - 1) + 1 \leq e(P, F) \leq s(t - 1) + 2s$ , implying  $(t - 1)(s - 2) - 1 \leq 0$ , a contradiction. If  $t$  is even, Then  $2t(s - 1) + 1 \leq e(P, F) \leq ts$ , which implies  $t(s - 2) + 1 \leq 0$ , a contradiction again.

**Lemma 2.13** *Let  $P = x_1 x_2 x_3$  and  $Q = y_1 y_2 y_3$  be two disjoint paths of  $G$  with  $x_1 \in V_1$  and  $y_1 \in V_2$ . Let  $F = (U_1, U_2; W)$  be a forest in  $G$  with  $|U_1| = |U_2| = s$ , where  $s \geq 3$ . Suppose  $F$  is disjoint from both  $P$  and  $Q$ , and  $d(x_1, F) + d(x_3, F) + d(y_1, F) + d(y_3, F) \geq 4s - 2$ . Then  $G[V(F \cup P \cup Q)] \supseteq F \dot{\cup} C$ , where  $C$  is a 4-cycle.*

**Proof.** First we claim that  $N(x_1, F) \cap N(x_3, F) \neq \emptyset$  and  $N(y_1, F) \cap N(y_3, F) \neq \emptyset$ . Suppose not, without loss of generality, say  $N(x_1, F) \cap N(x_3, F) = \emptyset$ , then  $d(x_1, F) + d(x_3, F) \leq s$ . It follows that  $2s \geq d(y_1, F) + d(y_3, F) \geq 4s - 2 - s = 3s - 2$ , implying  $s \leq 2$ , a contradiction. Clearly there exists one of  $\{x_1, x_3, y_1, y_3\}$ , say  $x_1$ , such that  $d(x_1, F) = s$ . Let  $u \in N(y_1, F) \cap N(y_3, F)$ . Then we see that  $F - u + x_1 \supseteq F$  and  $uQu$  is a 4-cycle disjoint from  $F - u + x_1$ , where  $u \in N(y_1, F) \cap N(y_3, F)$ .

### 3 Proofs of the Theorems

To prove the theorems, we introduce the following terminology: For a graph  $H$  and a path  $P = x_1x_2\dots x_t$  of  $H$ , we define  $\sigma(P, H) = \max\{d(x_2, H), d(x_{t-1}, H)\}$  if  $t \geq 2$  and  $\sigma(P, H) = d(x_1, H)$  if  $t = 1$ .

Let  $G$  and  $F$  be given as stated in the two theorems. We may assume that  $F$  is connected. If  $s = 2$ ,  $F$  is a path of order 4. Since  $|V_1| = |V_2| = n \geq 2k + 2 = 2(k + 1)$  and  $\delta(G) \geq k + 2 > k + 1$ , we see that if  $s = 2$  then  $G \supseteq F \dot{\cup} \bigcirc^k$  by Theorem B and Theorem C. Therefore we suppose  $s \geq 3$  and need to show the following:

$$\begin{aligned} G &\supseteq \bigcirc^k \dot{\cup} F \text{ if } n > 2k + s \text{ and} \\ G &\supseteq \bigcirc^{k-1} \dot{\cup} F \dot{\cup} P \text{ if } n = 2k + s, \text{ where } P \text{ is a path of order 4.} \end{aligned} \quad (1)$$

We use induction on  $k$  to prove (1). If  $k = 0$ , (1) follows from Lemma 2.9. Since  $n \geq 2k + s = 2(k - 1) + (s + 2)$  and  $\delta(G) \geq k + s = (k - 1) + (s + 1)$ , by induction on  $k$ ,  $G \supseteq \bigcirc^{k-1} \dot{\cup} F \dot{\cup} K_2$ . Let  $C_1, C_2, \dots, C_{k-1}$  be  $k - 1$  disjoint cycles of  $G$ . Let  $\sigma$  be an embedding of  $F$  in  $G - V(\bigcup_{i=1}^{k-1} C_i)$ . We choose  $C_1, C_2, \dots, C_{k-1}$  and  $\sigma(F)$  such that

$$\sum_{i=1}^{k-1} l(C_i) \quad \text{is minimum.} \quad (2)$$

Subject to (2), we choose  $C_1, C_2, \dots, C_{k-1}$  and  $\sigma(F)$  such that

$$e(G[\sigma(F)]) \quad \text{is maximum.} \quad (3)$$

Let  $D = G - V(\bigcup_{i=1}^{k-1} C_i) - V(\sigma(F))$ . Subject to (2) and (3), we choose  $C_1, C_2, \dots, C_{k-1}$  and  $\sigma(F)$  such that

$$\text{the length of a longest path in } D \text{ is maximal.} \quad (4)$$

Let  $P = x_1x_2\dots x_p$  be a fixed longest path of  $D$ . Without loss of generality, assume  $x_1 \in V_1$ . Subject to (2), (3) and (4), we choose  $C_1, C_2, \dots, C_{k-1}$  and  $\sigma(F)$  such that

$$\sigma(P, D) \quad \text{is minimum.} \quad (5)$$

Let  $D_0 = D - V(P)$ . Subject to (2) to (5), we choose  $C_1, C_2, \dots, C_{k-1}$  and  $\sigma(F)$  such that

$$\text{the length of a longest path in } D_0 \text{ is maximal.} \quad (6)$$

Let  $Q = y_1 y_2 \dots y_q$  be a fixed longest path of  $D_0$ . Without loss of generality, assume  $y_1 \in V_1$  if  $q$  is even. Subject to (2) to (6), we finally choose  $C_1, C_2, \dots, C_{k-1}$  and  $\sigma(F)$  such that

$$\text{if } q \text{ is odd, then } \sigma(Q, D_0) \text{ is minimum;} \quad (7)$$

$$\text{if } q \text{ is even, then } d(y_2, D_0) \text{ is minimum.} \quad (8)$$

Clearly,  $p \geq q$ . Let  $H = \bigcup_{i=1}^{k-1} C_i$  and  $|V(D)| = 2d$ . We will prove a number of claims. First, we claim

$$d \geq 2. \quad (9)$$

*Proof of (9).* Suppose  $d \leq 1$ . Without loss of generality, assume that  $l(C_1) \leq l(C_2) \leq \dots \leq l(C_{k-1})$  and  $l(C_{k-1}) = 2t$ . Then  $t \geq 3$ , for otherwise  $n = 2(k-1) + s + 1 < 2k + s$ . By Lemma 2.8 and (2),  $e(C_{k-1}, C_i) \leq 2t$  for all  $i \in \{1, \dots, k-2\}$ . By Lemma 2.1 and (2),  $d(x, C_{k-1}) \leq 1$  for all  $x \in V(D)$ . Therefore  $e(C_{k-1}, \sigma(F)) \geq 2t(k+s) - 2t(k-2) - 4t - 2 = 2ts - 2$ . Then  $G[V(C_{k-1} \cup \sigma(F))] \supseteq C' \dot{\cup} F$  by Lemma 2.10, where  $C'$  is a 4-cycle, contradicting (2).

We claim

$$p \geq 3 \text{ and if } |V(D_0)| \geq 4 \text{ then } q \geq 3. \quad (10)$$

*Proof of (10).* First we show  $p \geq 3$ . To the contrary, suppose  $p \leq 2$ . If  $p < 2$ , then for any  $x \in V(D) \cap V_1$  and  $y \in V(D) \cap V_2$ ,  $d(x, D) = d(y, D) = 0$ . It follows that  $d(x, H) + d(y, H) \geq 2(k+s) - 2s = 2k$ . Then there exists a  $C_i$  in  $H$  such that  $d(x, C_i) + d(y, C_i) \geq 3$ . By Lemma 2.1 and (2),  $C_i$  is a 4-cycle. By Lemma 2.2,  $G[V(C_i) \cup \{x, y\}] \supseteq C'_i \dot{\cup} K_2$ , where  $C'_i$  is a 4-cycle. This is a contradiction to  $p < 2$ . So  $p = 2$ . Let  $P = x_1 x_2$ . We may choose  $C_1, C_2, \dots, C_{k-1}$  and  $\sigma(F)$  such that  $D_0 \supseteq K_2$  while (2), (3) and (4) are maintained. If this is not the case, then by (4),  $d(x, D) = 0$  for all  $x \in D_0$ . For any  $x \in V(D_0) \cap V_1$  and  $y \in V(D_0) \cap V_2$ , if there exists a cycle, say  $C_1$ , such that  $d(x, C_1) + d(y, C_1) \geq 3$ , then by Lemma 2.1 and (2),  $C_1$  must be a 4-cycle. By Lemma 2.2,  $G[V(C_1) \cup \{x, y\}]$  contains a 4-cycle  $C'$  and an edge  $e'$  disjoint from  $C'$ . So we may assume  $d(x, C_i) + d(y, C_i) \leq 2$  for all  $C_i \in H$ . It follows that  $d(x, \sigma(F)) + d(y, \sigma(F)) \geq 2(k+s) - 2(k-1) = 2s + 2$ , a contradiction. Hence  $D_0 \supseteq K_2$ . This argument allows us to choose  $C_1, C_2, \dots, C_{k-1}$  and  $\sigma(F)$  such that  $D$  has a perfect matching. Let  $uv \in E(D_0)$  and  $R = \{x_1, x_2, u, v\}$ . If there exists a cycle  $C_i$  in  $H$  such that  $\sum_{x \in R} d(x, C_i) \geq 5$ , then by Lemma 2.5,  $G[V(C_i) \cup R]$  contains the disjoint union of a 4-cycle and a path of order 4, contradicting  $p = 2$ . So  $\sum_{x \in R} d(x, C_i) \leq 4$  for all  $C_i \in H$ . Therefore  $\sum_{x \in R} d(x, \sigma(F)) \geq 4(k+s) - 4(k-1) - 4 = 4s$ , i.e.  $d(x, \sigma(F)) = s$  for all  $x \in R$ . Clearly  $G[V(\sigma(F)) \cup R] \supseteq F \dot{\cup} C$ , where  $C$  is a 4-cycle, implying (1). Hence  $p \geq 3$ .

Suppose  $q \leq 2$  when  $|V(D_0)| \geq 4$ . By a similar argument, we may choose  $C_1, C_2, \dots, C_{k-1}$ ,  $\sigma(F)$  and  $P$  such that  $D_0 \supseteq 2K_2$ . Let  $u_1 v_1$  and  $u_2 v_2$  be two independent edges in  $D_0$ , and  $T = \{u_1, v_1, u_2, v_2\}$ . Since  $D$  is acyclic,  $\sum_{x \in T} d(x, D) \leq 6$ . By Lemmas 2.1 and 2.5,  $\sum_{x \in T} d(x, C_i) \leq 4$  for all  $C_i \in H$ . So  $\sum_{x \in T} d(x, \sigma(F)) \geq 4(k +$

$s) - 4(k - 1) - 6 = 4s - 2$ . Clearly there exists  $x \in T$  such that  $d(x, \sigma(F)) = s$ . Then  $G[V(\sigma(F))] = K_{s,s}$  follows from (3). By Lemma 2.11,  $G[V(\sigma(F)) \cup T] \supseteq F \dot{\cup} Q'$ , where  $Q'$  is a path of order 4 while (2), (3), (4), (5) are maintained, contradicting  $q \leq 2$ . Hence (10) holds.

The argument in the above paragraph shows that if  $|V(D_0)| \geq 2$ , then  $q \geq 2$ . We claim

$$\sigma(P, D) = 2, \sigma(Q, D_0) \leq 2 \text{ if } q \text{ is odd and } d(y_2, D_0) \leq 2 \text{ if } q \text{ is even.} \quad (11)$$

*Proof of (11).* First we suppose that  $\sigma(Q, D_0) \geq 3$  if  $q$  is odd and  $d(y_2, D_0) \geq 3$  if  $q$  is even. In the former case, we may assume  $d(y_2, D_0) \geq 3$  and  $q \geq 3$ . Let  $\{a, b\} = \{1, 2\}$  such that  $y_1 \in V_a$ . Let  $u$  be an endvertex of  $D_0$  such that  $uy_2 \in E$  and  $u \notin \{y_1, y_q\}$ . Clearly, either  $d(u, P) = 0$  or  $d(y_1, P) = 0$  as  $D$  is acyclic. Without loss of generality, assume that  $d(u, P) = 0$ . Let  $(A, B)$  be the bipartition of  $D_0 - V(Q) \cup \{u\}$  with  $A \subseteq V_a$  and  $B \subseteq V_b$ . Clearly  $|B| > |A|$ , so  $D_0 - V(Q) \cup \{u\}$  has a component  $T$  such that  $|V(T) \cap B| > |V(T) \cap A|$ . As there is at most one edge between  $Q$  and  $T$  and by Lemma 2.3, we can choose a vertex  $v \in V(T) \cap B$  such that  $d(v, D_0) \leq 1$ . We deduce that  $d(u, D) + d(v, D) \leq 3$  as  $D$  is acyclic.

If there exists  $C_i$  in  $H$  such that  $d(u, C_i) + d(v, C_i) \geq 3$ , then by Lemma 2.1 and (2),  $C_i$  must be a 4-cycle. By Lemma 2.2,  $G[V(C_i) \cup \{u, v\}] \supseteq C' \dot{\cup} e'$ , where  $C'$  is a 4-cycle and  $e'$  is an edge, and exactly one of  $u$  and  $v$  is an endvertex of  $e'$ . Let  $D' = G - (V(\bigcup_{j \neq i} C_j) \cup V(C')) - V(\sigma(F))$  and  $D'_0 = D' - V(P)$ . By (4),  $P$  is still a longest path of  $D'$ . So neither of the two endvertices of  $e'$  is adjacent to  $x_2$  or  $x_{p-1}$  and therefore  $\sigma(P, D') \leq \sigma(P, D)$ . Subsequently,  $Q$  is still a longest path of  $D'_0$  by (6). So neither of the two endvertices of  $e'$  is adjacent to  $y_2$  or  $y_{q-1}$ . Thus  $u \in V(C')$ ,  $d(y_2, D'_0) = d(y_2, D_0) - 1$  and  $d(y_{q-1}, D'_0) \leq d(y_{q-1}, D_0)$ . Repeating this argument for  $y_{q-1}$  if  $q$  is odd and  $d(y_{q-1}, D'_0) \geq 3$ , we obtain a contradiction with (7) or (8) while (2) to (6) are maintained.

So we may assume  $d(u, C_i) + d(v, C_i) \leq 2$  for all  $C_i \in H$ . It follows that  $d(u, \sigma(F)) + d(v, \sigma(F)) \geq 2(k + s) - 2(k - 1) - 3 = 2s - 1$ . By (3), it is easy to see that  $G[V(\sigma(F))] = K_{s,s}$ . If  $d(v, \sigma(F)) = s$ , then  $d(u, \sigma(F)) \geq s - 1$ . Clearly  $G[V(\sigma(F) \cup D_0)] \supseteq K_{s,s} \dot{\cup} Q'$ , where  $Q'$  is a path with  $l(Q') > l(Q)$  without violating (2) to (5). Therefore  $d(u, \sigma(F)) = s$  and  $d(v, \sigma(F)) = s - 1$ . Let  $F' = \sigma(F) - w + u$  and  $D'_0 = D_0 - u + w$ , where  $w \in N(v, \sigma(F))$ . Then  $d(y_2, D'_0) = d(y_2, D_0) - 1$  and  $d(y_{q-1}, D'_0) \leq d(y_{q-1}, D_0)$ . If  $q$  is even, we obtain a contradiction to (8) while (2) to (6) are maintained. If  $q$  is odd, we can obtain a contradiction to (7) by applying the same argument to  $y_{q-1}$ . A similar but simpler argument shows that  $\sigma(P, D) = 2$  as we have no concerns for the priorities (6) to (8). So (11) holds.

We claim

$$p \geq 2d - 1 \quad (12)$$

*Proof of (12).* Suppose  $p \leq 2d - 2$ . We distinguish two cases:  $p$  is even or odd.

*Case 1.*  $p$  is even.

By (10),  $p \geq 4$ . Let  $R = \{x_1, x_p, y_1, y_2\}$ . By (11),  $d(y_1, D_0) + d(y_2, D_0) \leq 3$ . Since  $e(P, Q) \leq 1$  and  $d(x_1, D) + d(x_p, D) = 2$ ,  $\sum_{x \in R} d(x, D) \leq 6$ .

If there exists  $C_i$  in  $H$  such that  $\sum_{x \in R} d(x, C_i) \geq 5$ , then by Lemma 2.1 and (2),  $C_i$  must be a 4-cycle. Let  $C_i = u_1 u_2 u_3 u_4 u_1$ . Without loss of generality, assume  $\{u_1, x_1, y_1\} \subseteq V_1$ . Clearly, either  $d(x_1, C_i) + d(y_2, C_i) \geq 3$  or  $d(x_p, C_i) + d(y_1, C_i) \geq 3$ . Without loss of generality, say the former holds. By Lemma 2.2,  $G[V(C_i) \cup \{x_1, y_2\}]$  contains a 4-cycle  $C'$  and an edge  $e'$  disjoint from  $C'$  such that exactly one of  $x_1$  and  $y_2$  is an endvertex of  $e'$ . By (4),  $x_1$  is not an endvertex of  $e'$ . So  $d(x_1, C_i) = 2$  and  $d(y_2, C_i) = 1$ . As  $d(y_1, C_i) + d(x_p, C_i) \geq 2$ , we have either  $d(y_1, C_i) > 0$  or  $N(x_p, C_i) \cap N(y_2, C_i) \neq \emptyset$ . In either case, it is easy to see that  $G[V(C_i \cup P) \cup \{y_1, y_2\}] \supseteq C'' \cup P'$ , where  $C''$  is a 4-cycle and  $P'$  is a path of order  $p + 2$ , contradicting (4).

So we may assume  $\sum_{x \in R} d(x, C_i) \leq 4$  for all  $C_i \in H$ . It follows that

$$\sum_{x \in R} d(x, \sigma(F)) \geq 4(k + s) - 4(k - 1) - 6 = 4s - 2.$$

Clearly there exists  $z \in R$  such that  $d(z, \sigma(F)) = s$ , so  $G[V(\sigma(F))] = K_{s,s}$  by (3). we have either  $d(x_1, \sigma(F)) + d(y_2, \sigma(F)) \geq 2s - 1$  or  $d(x_p, \sigma(F)) + d(y_1, \sigma(F)) \geq 2s - 1$ . Without loss of generality, say the former holds. If  $d(y_2, \sigma(F)) = s$ , then we readily see that  $G[V(\sigma(F) \cup P) \cup \{y_1, y_2\}]$  contains  $K_{s,s}$  and a path of order  $p + 1$  which is disjoint from  $K_{s,s}$ , contradicting (4). So  $d(y_2, \sigma(F)) = s - 1$  and  $d(x_1, \sigma(F)) = s$ . And moreover,  $N(y_2, \sigma(F)) \cap N(x_p, \sigma(F)) = \emptyset$ , for otherwise  $G[V(\sigma(F) \cup D)] \supseteq K_{s,s} \cup P'$ , where  $P'$  is a path of order  $p + 2$ , contradicting (4). Therefore  $d(y_2, \sigma(F)) + d(x_p, \sigma(F)) \leq s$ . It follows that  $2s \geq d(y_1, \sigma(F)) + d(x_1, \sigma(F)) \geq 4s - 2 - s = 3s - 2$ , implying  $s \leq 2$ , a contradiction.

*Case 2.*  $p$  is odd.

Notice that  $|V(D_0)|$  is odd. We claim that if  $q = 3$ , then we may choose  $Q$  such that  $y_1 \in V_2$ . Suppose that this is not true, i.e.  $y_1 \in V_1$ . Let  $(A, B)$  be the bipartition of  $D_0 - V(Q)$  such that  $A \subseteq V_1$  and  $B \subseteq V_2$ . Then  $|B| = |A| + 2$ . As  $D$  is acyclic and by Lemma 2.3, we can choose a vertex  $y_0 \in B$  such that  $d(y_0, D_0) \leq 1$ . Clearly,  $d(y_0, P) \leq 1$  and  $d(y_1, P) + d(y_3, P) \leq 1$ . We may assume  $d(y_1, P) = 0$ . So  $d(y_0, D) + d(y_1, D) \leq 3$ .

If there exists a  $C_i$  in  $H$  such that  $d(y_0, C_i) + d(y_1, C_i) \geq 3$ , then by Lemma 2.1, (2) and Lemma 2.2,  $C_i$  must be a 4-cycle, and moreover,  $G[V(C_i) \cup \{y_0, y_1\}]$  contains a 4-cycle  $C'$  and an edge  $e'$  disjoint from  $C'$  such that exactly one of  $y_0$  and  $y_1$  is an endvertex of  $e'$ . Replacing  $C_i$  with  $C'$  and by (4), we see that neither of the two endvertices of  $e'$  is adjacent to a vertex in  $\{x_1, x_2, x_{p-1}, x_p\}$ . Therefore (2) to (5) are maintained. By (6),  $y_1$  is not an endvertex of  $e'$ . So  $e' = y_0 z_0$  for some  $z_0 \in V(C_i)$ . Let  $H' = (H - V(C_i)) \cup C'$ ,  $D' = D - y_1 + z_0$  and  $D'_0 = D' - V(P)$ . Then  $D'_0$  does not contain a path of order 3 with its two endvertices in  $V_2$ . It follows from (11) that  $d(y_2, D'_0) = 1$ . Furthermore,  $\sum_{z \in S} d(z, D'_0) \leq 5$ , where  $S = \{y_2, y_3, y_0, z_0\}$ . As  $D'$  is acyclic,  $\sum_{z \in S} d(z, D') \leq 7$ . We distinguish two subcases:

Subcase 1.1. There exists a cycle  $C''$  in  $H'$  such that  $\sum_{z \in S} d(z, C'') \geq 5$ .



By Lemma 2.1 and (2),  $C'''$  must be a 4-cycle. By Lemma 2.5,  $G[V(C''') \cup S]$  contains a 4-cycle  $C'''$  and a path  $Q'$  of order 4 such that  $Q'$  is disjoint from  $C'''$ . By (4), no vertex of  $Q'$  is adjacent to a vertex in  $\{x_1, x_2, x_{p-1}, x_p\}$ . Thus we obtain a contradiction to (6) while (2) to (5) are maintained.

Subcase 1.2.  $\sum_{z \in S} d(z, C'_i) \leq 4$  for all  $C'_i \in H'$ .

Clearly  $\sum_{z \in S} d(z, \sigma(F)) \geq 4(k+s) - 7 - 4(k-1) = 4s - 3$ . Then there exists  $z \in S$  such that  $d(z, \sigma(F)) = s$ . It follows from (3) that  $G[V(\sigma(F))] = K_{s,s}$ . By Lemma 2.11,  $G[V(\sigma(F) \cup Q) \cup \{y_0, z_0\}] \supseteq F \dot{\cup} Q'$ , where  $Q'$  is a path of order 4, contradicting  $q = 3$ .

So we may assume  $d(y_0, C_i) + d(y_1, C_i) \leq 2$  for all  $C_i \in H$ . Consequently

$$d(y_0, \sigma(F)) + d(y_1, \sigma(F)) \geq 2(k+s) - 2(k-1) - 3 = 2s - 1.$$

If  $d(y_0, \sigma(F)) = s$ , it's easy to see that  $G[V(\sigma(F)) \cup \{y_1, y_2, y_3, y_0\}]$  contains  $F$  and a disjoint path of order 4, contradicting  $q = 3$ . So  $d(y_0, \sigma(F)) = s - 1$  and  $d(y_1, \sigma(F)) = s$ . Let  $y_0 z_0 \in E$  for some  $z_0 \in V(\sigma(F))$ . By (6),  $y_2 z_0 \notin E$ . Let  $\sigma'(F) = \sigma(F) - z_0 + y_1$ ,  $D'_0 = D_0 - y_1 + z_0$  and  $D' = D'_0 \cup P$ . Then  $d(y_2, D'_0) = 1$ , and moreover,  $d(z_0, D'_0) \leq 1$  for otherwise we have a path of order 3 with both endvertices in  $V_2$ . Let  $T = \{y_2, y_3, y_0, z_0\}$ . Then  $\sum_{z \in T} d(z, D) \leq 7$  as  $\sum_{z \in T} d(z, P) \leq 2$ . Therefore  $\sum_{z \in T} d(z, \sigma'(F)) \geq 4(k+s) - 4(k-1) - 7 = 4s - 3$ . Again  $G[V(\sigma'(F))] = K_{s,s}$  follows from (3). By Lemma 2.11,  $G[V(\sigma'(F)) \cup T] \supseteq F \dot{\cup} Q'$ , where  $Q'$  is a path of order 4, contradicting  $q = 3$ .

Now  $y_1 \in V_2$  for  $q = 3$ , so we can choose three distinct vertices  $z_1, z_2, z_3$  from  $D_0$  with  $z_1 \in V_1$  and  $\{z_2, z_3\} \subseteq V_2$  such that  $\{z_1, z_2\} = \{y_1, y_2\}$ , and if  $q \geq 3$  then  $z_3 \in \{y_{q-1}, y_q\}$ . If  $q = 2$ , then  $|V(D_0)| = 3$  by (10) and therefore  $z_3$  is an isolated vertex of  $D_0$ . Let  $T = \{x_1, x_{p-1}, x_p, z_1, z_2, z_3\}$ . As  $D$  is acyclic and  $d(z_3, P) \leq 1$ , we deduce from (11) that  $\sum_{u \in T} d(u, D) \leq 10$ .

If there exists a  $C_i$  in  $H$  such that  $\sum_{u \in T} d(u, C_i) \geq 7$ , then by Lemma 2.1 and (2),  $C_i$  must be a 4-cycle. Let  $C_i = v_1 v_2 v_3 v_4 v_1$  with  $v_1 \in V_1$ . If  $d(z_2, C_i) = 2$  or  $d(z_3, C_i) = 2$ , it is easy to see, by observing two situations that either  $d(x_1, C_i) + d(x_p, C_i) \geq 1$  or  $d(x_1, C_i) + d(x_p, C_i) = 0$ , that  $G[V(C_i \cup P) \cup \{z_1, z_2, z_3\}]$  contains a 4-cycle  $C'$  and a path  $P'$  disjoint from  $C'$  but longer than  $P$ , contradicting (4). Hence  $d(z_2, C_i) \leq 1$  and  $d(z_3, C_i) \leq 1$ . We distinguish two subcases. Note that  $z_1 z_2 \in E$ .

Subcase 2.1.  $q \geq 3$ .

We first suppose that  $d(z_1, C_i) \geq 1$  and  $d(z_2, C_i) = 1$ . Without loss of generality, say  $\{v_1 z_2, v_2 z_1\} \subseteq E$ . Then  $C' = v_1 v_2 z_1 z_2 v_1$  is a 4-cycle, and  $e(\{x_1, x_{p-1}, x_p\}, \{v_3, v_4\}) = 0$  By (4). As  $\sum_{u \in T} d(u, C_i) \geq 7$ , we deduce that  $d(u, C_i) = 1$  for all  $u \in T - \{z_1\}$  and  $d(z_1, C_i) = 2$ . Then  $z_1 z_2 v_1 v_4 z_1$  and  $v_2 P v_2$  are two disjoint cycles in  $G[V(C_i \cup P) \cup \{z_1, z_2\}]$ . So either  $d(z_1, C_i) = 0$  or  $d(z_2, C_i) = 0$ . Suppose the former holds. We have  $d(x_1, C_i) + d(x_{p-1}, C_i) + d(x_p, C_i) \geq 5$  and therefore  $N(x_1, C_i) \cap N(x_p, C_i) \neq \emptyset$ . For  $v_2 \in N(x_1, C_i) \cap N(x_p, C_i)$ , clearly  $G[V(C_i \cup Q)] - v_2$  is disjoint from  $v_2 P v_2$  and therefore is acyclic. So  $d(z_2, C_i) + d(z_3, C_i) \leq 1$ . Consequently,  $d(x_1, C_i) =$

$d(x_{p-1}, C_i) = d(x_p, C_i) = 2$  and  $d(z_j, C_i) = 1$  for some  $j \in \{2, 3\}$ . Without loss of generality, say  $z_j v_1 \in E$ . Then the 4-cycle  $x_{p-1} x_p v_4 v_3 x_{p-1}$  is disjoint from the path  $z_j v_1 v_2 x_1 x_2 \dots x_{p-2}$  which is longer than  $P$ , contradicting (4). Therefore  $d(z_1, C_i) > 0$  and  $d(z_2, C_i) = 0$ .

If  $d(z_3, C_i) = 0$ , then there exists  $u' \in \{x_1, x_{p-1}, x_p, z_1\}$  such that  $d(u, C_i) = 2$  for all  $u \in \{x_1, x_{p-1}, x_p, z_1\} - \{u'\}$  and  $d(u', C_i) \geq 1$ . This implies that  $\{v_i z_1, v_i x_1, v_j x_p\} \subseteq E$  for some  $\{i, j\} = \{2, 4\}$  and  $x_{p-1} v_h \in E$  for some  $h \in \{1, 3\}$ . Then the 4-cycle  $x_{p-1} x_p v_j v_h x_{p-1}$  is disjoint from the path  $z_2 z_1 v_i x_1 x_2 \dots x_{p-2}$  which is longer than  $P$ , contradicting (4). Therefore  $d(z_3, C_i) = 1$ . Say  $\{v_1 z_3, v_2 z_1\} \subseteq E$ . Then  $G[V(Q) \cup \{v_1, v_2\}]$  contains a cycle and therefore  $G[V(P) \cup \{v_3, v_4\}]$  is acyclic. Hence

$$e(\{x_1, x_{p-1}, x_p\}, \{v_3, v_4\}) \leq 1.$$

This implies that  $d(x_1, C_i) + d(x_{p-1}, C_i) + d(x_p, C_i) = 4$  as  $d(z_1, C_i) + d(z_3, C_i) \leq 3$ . Thus  $d(z_1, C_i) = 2$  and  $x_{p-1} v_1 \in E$ . Then the 4-cycle  $z_1 v_2 v_3 v_4 z_1$  is disjoint from the path  $x_1 x_2 \dots x_{p-1} v_1 z_3$  which is longer than  $P$ , contradicting (4) again.

Subcase 2.2.  $q = 2$ . Notice that  $d(z_3, D) \leq 1$ .

First suppose that there exists  $C_i$  in  $H$  such that  $d(x_p, C_i) + d(z_3, C_i) \geq 3$ , then by Lemma 2.1, Lemma 2.2, (2) and (3) as before, we see that  $C_i$  is a 4-cycle,  $d(x_p, C_i) = 2$  and  $d(z_3, C_i) = 1$ . Let  $L_1 = C_i - z_4 + x_p$  where  $z_4 \in V(C_i)$  such that  $z_3 z_4 \in E$ . Let  $H_1 = (H - V(C_i)) \cup L_1$  and  $D_1 = G - V(H_1) - V(\sigma(F))$ . As  $D_1$  is acyclic,  $\sum_{i=1}^4 d(z_i, D_1) \leq 7$ . If there exists a cycle  $C'$  in  $H_1$  such that  $\sum_{i=1}^4 d(z_i, C') \geq 5$ , then by Lemma 2.1 and (2),  $C'$  must be a 4-cycle. By Lemma 2.5,  $G[V(C') \cup \{z_1, z_2, z_3, z_4\}] \supseteq C'' \dot{\cup} Q'$ , where  $C''$  is a 4-cycle and  $Q'$  is a path of order 4. If  $\sum_{i=1}^4 d(z_i, C'_i) \leq 4$  for all  $C'_i \in H_1$ , then  $\sum_{i=1}^4 d(z_i, \sigma(F)) \geq 4(k+s) - 7 - 4(k-1) = 4s - 3$ . Again  $G[V(\sigma(F))] = K_{s,s}$  by (3). It follows from Lemma 2.11 that  $G[V(\sigma(F)) \cup \{z_1, z_2, z_3, z_4\}] \supseteq F \dot{\cup} Q'$ , where  $Q'$  is a path of order 4. So in both cases we obtain a path  $Q'$  of order 4. Without loss of generality, say the former case holds. As  $p$  is odd and by (4),  $p \geq 5$ . Let  $H_2 = (H_1 - V(C')) \cup C''$ ,  $D_2 = G - V(H_2) - V(\sigma(F))$ ,  $P' = P - x_p$  and  $Q' = u_1 u_2 u_3 u_4$  with  $u_1 \in V_1$ . Then  $D_2$  is acyclic and  $e(P', Q') \leq 1$ .

When  $p \geq 7$ , if there exists a cycle  $C'''$  in  $H_2$  such that  $\sum_{i=1}^{p-1} d(x_i, C''') \geq p$ , then by Lemma 2.1 and (2),  $C'''$  must be a 4-cycle. It follows from Lemma 2.7 that  $G[V(C''' \cup P')] \supseteq \bigcirc^2$ , implying (1). So we may assume  $\sum_{i=1}^{p-1} d(x_i, C''_i) \leq p-1$  for all  $C''_i \in H_2$ . Therefore  $\sum_{i=1}^{p-1} d(x_i, \sigma(F)) \geq (p-1)(k+s) - 2(p-2) - 1 - (p-1)(k-1) = (s-1)(p-1) + 1$ . By Lemma 2.12,  $G[V(\sigma(F) \cup P)] \supseteq F \dot{\cup} \bigcirc$ , which implies (1).

When  $p = 5$ , we have  $e(\{x_1, x_3\}, \{u_2, u_4\}) = 0$ . Let  $W = \{x_1, x_3, u_2, u_4\}$ . Then  $\sum_{w \in W} d(w, D_2) = 6$  as  $D_2$  is acyclic. If there exists a cycle  $L'$  in  $H_2$  such that  $\sum_{w \in W} d(w, L') \geq 5$ , then by Lemma 2.1 and (2),  $L'$  must be a 4-cycle. By Lemma 2.4,  $G[V(L') \cup \{x_1, x_2, x_3, u_2, u_3, u_4\}] \supseteq L'' \dot{\cup} P''$ , where  $L''$  is a 4-cycle and  $P''$  is a path of order 6, contradicting  $p = 5$ . So  $\sum_{w \in W} d(w, L_i) \leq 4$  for all  $L_i \in H_2$ . Therefore  $\sum_{w \in W} d(w, \sigma(F)) \geq 4(k+s) - 6 - 4(k-1) = 4s - 2$ . Evidently (1) follows from Lemma 2.13.

So we can assume  $d(x_p, C_i) + d(z_3, C_i) \leq 2$  for all  $C_i \in H$ , then  $d(x_p, \sigma(F)) + d(z_3, \sigma(F)) \geq 2(k+s) - 2 - 2(k-1) = 2s$ . Clearly  $G[V(\sigma(F) \cup P)] \supseteq F \dot{\cup} P'$ , where  $P'$  is a path of order  $p+1$ , a contradiction to (4). This proves the subcase 2.2.

Now we may assume that  $\sum_{u \in T} d(u, C_i) \leq 6$  for all  $C_i \in H$ . Then

$$\sum_{u \in T} d(u, \sigma(F)) \geq 6(k+s) - 10 - 6(k-1) = 6s - 4.$$

Again  $G[V(\sigma(F))] = K_{s,s}$  by (3). We claim that there exists  $x \in \{x_1, x_p\}$ , say  $x_1$ , such that  $d(x_1, \sigma(F)) \geq 1$ . Suppose that this is not the case, then  $d(x_1, \sigma(F)) = d(x_p, \sigma(F)) = 0$ . It follows that  $4s \geq d(x_{p-1}, \sigma(F)) + d(z_1, \sigma(F)) + d(z_2, \sigma(F)) + d(z_3, \sigma(F)) \geq 6s - 4$ , implying  $s \leq 2$ , a contradiction. Similarly there exists  $z \in \{z_2, z_3\}$  say  $z_2$  such that  $d(z_2, \sigma(F)) \geq 1$ . Let  $\{ux_1, vz_2\} \subseteq E$ , where  $\{u, v\} \subseteq V(\sigma(F))$ . Then  $\sigma(F) - u + z_2 \supseteq F$  and  $P + u$  is a path disjoint from  $F$ , a contradiction to (4). So (12) holds.

We are now in the position to complete the proofs. By (9) and (12),  $p \geq 2d - 1 \geq 3$ . As  $D$  is acyclic,  $e(P, D) \leq 2(p-1) + 1$ . We distinguish two cases:

*Case 1.* There exists a  $C_i$  in  $H$  such that  $e(P, C_i) \geq p + 1$ .

By Lemma 2.1 and (2),  $C_i$  must be a 4-cycle. If  $p \geq 6$ , then by Lemma 2.7,  $G[V(C_i \cup P)] \supseteq \bigcirc^2$ , implying (1). So assume  $p \leq 5$  and therefore  $d = 2$  or  $d = 3$ .

If  $d = 2$ , we will prove Theorem 2. First we prove  $p = 4$ . If  $p \neq 4$ , then by (10),  $p = 3$ . Without loss of generality, assume  $\{x_1, x_3\} \subseteq V_1$ . Let  $x_0 \in D - V(P)$ . Clearly  $d(x_0, D) + d(x_3, D) = 1$ . If there exists a cycle  $C_i$  in  $H$  such that  $d(x_3, C_i) + d(x_0, C_i) \geq 3$ , then by Lemma 2.1 and (2),  $C_i$  must be a 4-cycle and  $G[V(C_i) \cup \{x_0, x_3\}]$  contains a 4-cycle  $C'$  and an edge  $e'$  disjoint from  $C'$ , and moreover, we must have  $e' = x_0z$  for some  $z \in V(C_i)$ , for otherwise  $G[V(C_i \cup D)] \supseteq C'_i \dot{\cup} L$ , where  $C'_i$  is a 4-cycle and  $L$  is a path of order 4, a contradiction. Let  $D' = D - x_3 + z$  and  $H' = (H - V(C_i)) \cup C'$ . If there exists a cycle, say  $C'_1$  in  $H'$  such that  $e(D', C'_1) \geq 5$ , then by Lemma 2.5,  $G[V(C'_1 \cup D')]$  contains a 4-cycle and a disjoint path of order 4, contradicting  $p = 3$ . So we may assume  $e(D', C'_i) \leq 4$  for all  $C'_i \in H'$ . It follows that  $e(D', \sigma(F)) \geq 4(k+s) - 4(k-1) - 4 = 4s$ , which implies  $G[V(\sigma(F) \cup D')]$  contains a disjoint path of order 4, a contradiction. Thus  $d(x_3, C_i) + d(x_0, C_i) \leq 2$  for all  $C_i \in H$ , implying  $d(x_3, \sigma(F)) + d(x_0, \sigma(F)) \geq 2(k+s) - 1 - 2(k-1) = 2s + 1$ , a contradiction again. Hence  $p = 4$ .

Now we prove  $n = 2k + s$ . Suppose  $l(C_1) \leq l(C_2) \leq \dots \leq l(C_{k-1}) = 2t$ . It's enough to show  $t = 2$ . If  $t \geq 3$ , then by Lemma 2.8 and (2),  $e(C_{k-1}, C_i) \leq 2t$  for all  $i \in \{1, \dots, k-2\}$ , and moreover,  $e(C_{k-1}, P) \leq 4$  by Lemma 2.1 and (2). Therefore  $e(C_{k-1}, \sigma(F)) \geq 2t(k+s) - 2t(k-2) - 4t - 4 = 2ts - 4$ . By Lemma 2.10,  $G[V(C_{k-1} \cup \sigma(F))]$  contains a disjoint path of order 4, contradicting  $t \geq 3$ . Hence Theorem 2 holds.

If  $d = 3$ , then  $p = 5$ . Let  $z_0 \in V(D) - V(P)$ . If  $d(x_1, C_i) + d(z_0, C_i) \leq 2$  for all  $C_i \in H$ , then  $d(x_1, \sigma(F)) + d(z_0, \sigma(F)) \geq 2(k+s) - 2(k-1) - 2 = 2s$ . Clearly  $G[V(\sigma(F) \cup D)] \supseteq F \dot{\cup} L$ , where  $L$  is a path of order 6, a contradiction to (4). So we may assume that there exists  $C_i \in H$ , say  $C_1$  such that  $d(x_1, C_1) + d(z_0, C_1) \geq 3$ .

As before, by Lemma 2.1, Lemma 2.2, (2) and (3), we see that  $C_1$  is a 4-cycle,  $d(x_1, C_1) = 2$  and  $d(z_0, C_1) = 1$ . Let  $H_1 = H - V(C_1)$  and  $z_1 \in V(C_1)$  be such that  $z_1 z_0 \in E$ . Consider  $\{x_5, z_0\}$ .

If there exists  $C_j \in H_1$ , say  $C_2$  such that  $d(x_5, C_2) + d(z_0, C_2) \geq 3$ . Then  $C_2$  is a 4-cycle,  $d(x_5, C_2) = 2$  and  $d(z_0, C_2) = 1$ . Let  $z_2 \in V(C_2)$  be such that  $z_0 z_2 \in E$ . Let  $H' = (H - V(C_1 \cup C_2)) \cup (C_1 - z_1 + x_1) \cup (C_2 - z_2 + x_5)$ ,  $D' = G - V(H') - V(\sigma(F))$  and  $U = \{x_2, x_4, z_1, z_2\}$ . Clearly  $H'$  consists of  $k - 1$  disjoint cycles satisfying (2). Then  $d(u, D') = 1$  for all  $u \in U$ , for otherwise  $D'$  contains a path of order 6, contradicting (4). If there exists  $C' \in H'$  such that  $\sum_{u \in U} d(u, C') \geq 5$ , then by Lemma 2.1 and (2),  $C'$  is a 4-cycle. By Lemma 2.4,  $G[V(C' \cup D')] \supseteq C'' \dot{\cup} P'$ , where  $C''$  is a 4-cycle and  $P'$  is a path of order 6, a contradiction. So we may assume  $\sum_{u \in U} d(u, C'_i) \leq 4$  for all  $C'_i \in H'$ . Therefore  $\sum_{u \in U} d(u, \sigma(F)) \geq 4(k + s) - 4(k - 1) - 4 = 4s$ . It follows that  $G[V(\sigma(F) \cup D')] \supseteq F \dot{\cup} C'''$ , where  $C'''$  is a 4-cycle, implying (1).

So we may suppose that  $d(x_5, C_i) + d(z_0, C_i) \leq 2$  for all  $C_i \in H_1$ . It follows that  $d(x_5, \sigma(F)) + d(z_0, \sigma(F)) \geq 2(k + s) - 2(k - 2) - 5 = 2s - 1$ . If  $d(z_0, \sigma(F)) = s$ , clearly  $G[V(\sigma(F) \cup D)] \supseteq F \dot{\cup} L$ , where  $L$  is a path of order 6, contradicting  $p = 5$ . So we may assume  $d(z_0, \sigma(F)) = s - 1$  and  $d(x_5, \sigma(F)) = s$ . Let  $w \in N(z_0, \sigma(F))$  and  $W = \{x_2, x_4, z_1, w\}$ . It's easy to see that  $G[V(C_1 \cup D \cup \sigma(F))] \supseteq C'_1 \dot{\cup} D' \dot{\cup} F$ , where  $C'_1$  is a 4-cycle and  $D' = G[\{x_2, x_3, x_4, z_1, z_0, w\}]$ . If  $\sum_{u \in W} d(u, \sigma(F)) = 4s$ , then evidently  $G[V(\sigma(F) \cup D')] \supseteq F \dot{\cup} \bigcirc$ , implying (1). So we may assume  $e(W, \sigma(F)) \leq 4s - 1$ . Furthermore, we have  $e(W, D') = 4$ , thus  $e(W, H') \geq 4(k + s) - 4 - (4s - 1) = 4(k - 1) + 1$ , where  $H' = H_1 \cup C'_1$ . This implies that there exists a cycle  $C'$  in  $H'$  such that  $e(W, C') \geq 5$ . Again by Lemma 2.1 and (2),  $C'$  is a 4-cycle. By Lemma 2.4,  $G[V(C' \cup D')] \supseteq F \dot{\cup} P'$ , where  $P'$  is a path of order 6, a contradiction.

*Case 2.*  $e(P, C_i) \leq p$  for all  $C_i \in H$ .

We have  $e(P, \sigma(F)) \geq p(k + s) - p(k - 1) - (2(p - 1) + 1) = p(s - 1) + 1$ . If  $p$  is even, let  $p = 2t$ . If  $t = 2$  then  $d = 2$ . So assume  $t \geq 3$ . It follows from Lemma 2.12 that  $G[V(P \cup \sigma(F))] \supseteq F \dot{\cup} \bigcirc$ , implying (1). If  $p$  is odd, let  $p = 2t + 1$ . If  $t = 2$  then  $p = 5$ . So assume  $t \geq 3$ . If  $d(x_1, \sigma(F)) \leq s - 1$  or  $d(x_p, \sigma(F)) \leq s - 1$ , then let  $P' = P - x_1$  or  $P - x_p$ . We have  $e(P', \sigma(F)) \geq (2t + 1)(s - 1) + 1 - (s - 1) = 2t(s - 1) + 1$ . By Lemma 2.12,  $G[V(P' \cup \sigma(F))] \supseteq F \dot{\cup} \bigcirc$ . So  $d(x_1, \sigma(F)) = d(x_p, \sigma(F)) = s$ . Let  $T = \{x_{2i} : i = 1, \dots, (p - 1)/2\}$ . If there exists  $\{x, y\} \subseteq T$  such that  $N(x, \sigma(F)) \cap N(y, \sigma(F)) \neq \emptyset$ , then clearly  $G[V(P \cup \sigma(F))] \supseteq F \dot{\cup} \bigcirc$ . Therefore  $\sum_{x \in T} d(x, \sigma(F)) \leq s$ . Let  $U = \{x_{2i+1} : i = 1, \dots, (p - 3)/2\}$ . We have  $e(U, \sigma(F)) = 0$ , for otherwise  $G[V(P \cup \sigma(F))] \supseteq F \dot{\cup} \bigcirc$ . It follows that  $3s \geq e(P, \sigma(F)) \geq (2t + 1)(s - 1) + 1$ , implying  $(s - 1)(t - 1) \leq 1$ , a contradiction. This completes the proofs of the theorems.

## 4 References

- [1] B. Bollobás, *Extremal Graph Theory*, Academic Press, London (1978).
- [2] S. Brandt, Subtrees and subforests of graphs, *J. Combin. Theory, Ser. B* **61** (1994), 63–70.

- [3] K. Corrádi and A. Hajnal, On the maximal number of independent circuits in a graph, *Acta Math. Acad. Sci. Hungar.* **14** (1963), 423–439.
- [4] L. Lesniak, Independent cycles in graphs, *JCMCC* **17** (1995), 55–63.
- [5] G. W. Schuster, A minimum degree result for disjoint cycles and forests in graphs. *Combinatorica* **18**(3)(1998) 425–436.
- [6] H. Wang, Independent cycles with limited size in a graph. *Graphs and Combinatorics* **10** (1994), 271–281.
- [7] H. Wang, On the maximum number of independent cycles in a bipartite graph. *J. Combin. Theory, Ser. B* **67** (1996), 152–164.
- [8] H. Wang, On the maximum number of independent cycles in a graph. *Discrete Mathematics* **205** (1999), 183–190.

(Received 30 Jan 2002)