

New orthogonal designs from weighing matrices

ILIAS S. KOTSIREAS*

*Department of Physics and Computer Science
Wilfrid Laurier University
75 University Avenue West
Waterloo, Ontario N2L 3C5
Canada*

CHRISTOS KOUKOUVINOS

*Department of Mathematics
National Technical University of Athens
Zografou 15773, Athens
Greece*

JENNIFER SEBERRY

*Centre for Computer Security Research
School of Information Technology and Computer Science
University of Wollongong
Wollongong, NSW 2522
Australia*

Abstract

In this paper we show the existence of new orthogonal designs, based on a number of new weighing matrices of order $2n$ and weights $2n - 5$ and $2n - 9$ constructed from two circulants. These new weighing matrices were constructed recently by establishing various patterns on the locations of the zeros in a potential solution, in conjunction with the power spectral density criterion. We also demonstrate that some of these new orthogonal designs, the ones that are full, can be used to construct inequivalent Hadamard matrices.

1 Introduction

A weighing matrix $W = W(n, k)$ is a square matrix with entries $0, \pm 1$ having k non-zero entries per row and column and inner product of distinct rows equal to

* Supported by an NSERC grant.

zero. Therefore W satisfies $WW^t = kI_n$. The number k is called the weight of W . Weighing matrices have been studied extensively, see [1], [6], [2] and references therein, for detailed information on known and unknown weighing matrices.

A well-known necessary condition for the existence of $W(2n, k)$ matrices states that if there exists a $W(2n, k)$ matrix with n odd, then $k < 2n$ and k is the sum of two squares. In this paper we are focusing on $W(2n, k)$ constructed from two circulants. The two circulants construction for weighing matrices is described in the theorem below, taken from [3].

Theorem 1 *If there exist two circulant matrices A_1, A_2 of order n , with $0, \pm 1$ elements, satisfying $A_1A_1^t + A_2A_2^t = fI_n$ and f is an integer, then there exists a $W(2n, f)$, given as*

$$W(2n, f) = \begin{pmatrix} A_1 & A_2 \\ -A_2^t & A_1^t \end{pmatrix} \text{ or } W(2n, f) = \begin{pmatrix} A_1 & A_2R \\ -A_2R & A_1 \end{pmatrix}$$

where R is the square matrix of order n with $r_{ij} = 1$ if $i + j - 1 = n$ and 0 otherwise.

Recently, a number of new weighing matrices of order $2n$ and weights $2n - 5$, $2n - 9$ was constructed as per theorem 1 in [4, 5], by establishing patterns for the potential locations of zeros in the first rows of matrices A_1 and A_2 . The existence of these weighing matrices has certain implications on the existence of orthogonal designs and this is what we want to exhibit in this paper. It is also important to point out that the construction patterns used to find new weighing matrices in [4, 5], are of crucial important for the orthogonal design constructions of this paper.

The orthogonal design constructions of this paper are based on formulations of a theorem of Goethals and Seidel which may be used in the following form:

Theorem 2 [3, Theorem 4.49] *If there exist four circulant matrices A_1, A_2, A_3, A_4 of order n satisfying*

$$\sum_{i=1}^4 A_i A_i^T = fI$$

where f is the quadratic form $\sum_{j=1}^u s_j x_j^2$, then there is an orthogonal design $OD(4n; s_1, s_2, \dots, s_u)$.

2 New Orthogonal designs from $W(2n, 2n - 5)$

The first lemma is stated for completeness only.

Lemma 1 *If there exists an $OD(2n; a, b)$ constructed using two circulant matrices then, using the two circulant $W(2n, 2n - 5)$ constructed here with the Goethals-Seidel array we have $OD(4n; a, b, 2n - 5)$.*

The next theorem relies heavily on the construction pattern used to find the $W(2n, 2n - 5)$.

Theorem 3 *Suppose there exists a $W(2n, 2n - 5)$ constructed using two circulants with first rows of the required pattern. Then there exist*

1. $OD(4n; 4, 4, 4n - 10)$ and
2. $OD(4n; 2, 8, 4n - 10)$ constructed using four circulant matrices in the Goethals-Seidel array.

Proof We multiply each of the elements of the two $\{0, \pm 1\}$ sequences by the commuting variable a . We rearrange the required sequences of the $W(2n, 2n - 5)$, if necessary, to the form

$$\begin{aligned} A &= 0 \quad 0 \quad a_3 \quad \cdots \quad a_{\frac{n-1}{2}} \quad 0 \quad a_{\frac{n+3}{2}} \quad \cdots \quad a_n \\ B &= 0 \quad 0 \quad b_3 \quad \cdots \quad b_{\frac{n-1}{2}} \quad b_{\frac{n+1}{2}} \quad b_{\frac{n+3}{2}} \quad \cdots \quad b_n \end{aligned} \tag{1}$$

where each element a_i and b_j is $\pm a$.

Now let p, q, x and y be commuting variables.

Then

$$\begin{aligned} P &= p \quad q \quad a_3 \quad \cdots \quad a_{\frac{n-1}{2}} \quad 0 \quad a_{\frac{n+3}{2}} \quad \cdots \quad a_n \\ Q &= -p \quad -q \quad a_3 \quad \cdots \quad a_{\frac{n-1}{2}} \quad 0 \quad a_{\frac{n+3}{2}} \quad \cdots \quad a_n \\ R &= p \quad -q \quad b_3 \quad \cdots \quad b_{\frac{n-1}{2}} \quad b_{\frac{n+1}{2}} \quad b_{\frac{n+3}{2}} \quad \cdots \quad b_n \\ S &= -p \quad q \quad b_3 \quad \cdots \quad b_{\frac{n-1}{2}} \quad b_{\frac{n+1}{2}} \quad b_{\frac{n+3}{2}} \quad \cdots \quad b_n \end{aligned}$$

give the required sequences for the $OD(4n; 4, 4, 4n - 10)$ and

$$\begin{aligned} X &= y \quad -y \quad a_3 \quad \cdots \quad a_{\frac{n-1}{2}} \quad x \quad a_{\frac{n+3}{2}} \quad \cdots \quad a_n \\ Y &= -y \quad y \quad a_3 \quad \cdots \quad a_{\frac{n-1}{2}} \quad -x \quad a_{\frac{n+3}{2}} \quad \cdots \quad a_n \\ Z &= y \quad y \quad b_3 \quad \cdots \quad b_{\frac{n-1}{2}} \quad b_{\frac{n+1}{2}} \quad b_{\frac{n+3}{2}} \quad \cdots \quad b_n \\ W &= -y \quad -y \quad b_3 \quad \cdots \quad b_{\frac{n-1}{2}} \quad b_{\frac{n+1}{2}} \quad b_{\frac{n+3}{2}} \quad \cdots \quad b_n \end{aligned}$$

give the required sequences for the $OD(4n; 2, 8, 4n - 10)$.

Remark. We note that these last orthogonal designs are full and so give rise to Hadamard matrices.

3 New Orthogonal designs from $W(2n, 2n - 9)$

The first lemma is stated for completeness only.

Lemma 2 *If there exists an $OD(2n; a, b)$ constructed using two circulant matrices then, using the two circulant $W(2n, 2n - 9)$ constructed here with the Goethals-Seidel array we have $OD(4n; a, b, 2n - 9)$.*

The next theorem relies heavily on the construction pattern used to find the $W(2n, 2n - 9)$.

3.1 The 1-4-4 Pattern

Theorem 4 *Suppose there exists a $W(2n, 2n - 9)$ constructed using two circulants with first rows of the required pattern. Then there exist*

1. $OD(4n; 2, 4n - 18)$,
2. $OD(4n; 4, 4n - 18)$,
3. $OD(4n; 4, 4, 4n - 18)$,
4. $OD(4n; 2, 8, 4n - 18)$ and
5. $OD(4n; 8, 8, 4n - 18)$ constructed using four circulant matrices in the Goethals-Seidel array.

Proof We multiply each of the elements of the two $\{0, \pm 1\}$ sequences by the commuting variable a . We rearrange the required sequences of the $W(2n, 2n - 9)$, if necessary, to the form

$$\begin{aligned} A &= 0 \ 0 \ 0 \ 0 \ a_5 \ \cdots \ a_{\frac{n-1}{2}} \ 0 \ a_{\frac{n+3}{2}} \ \cdots \ a_n \\ B &= 0 \ 0 \ 0 \ 0 \ b_5 \ \cdots \ b_{\frac{n-1}{2}} \ b_{\frac{n+1}{2}} \ b_{\frac{n+3}{2}} \ \cdots \ b_n \end{aligned} \tag{2}$$

where each element a_i and b_j is $\pm a$.

Now let p, q, r, s, x, y, z and w be commuting variables. Cases 1) to 5) are constructed by forming the four circulant matrices with first rows which are then used in the Goethals-Seidel array

$$\begin{aligned} P &= p \quad q \quad r \quad s \quad a_5 \ \cdots \ a_{\frac{n-1}{2}} \quad 0 \quad a_{\frac{n+3}{2}} \ \cdots \ a_n \\ Q &= -p \quad -q \quad -r \quad -s \quad a_5 \ \cdots \ a_{\frac{n-1}{2}} \quad 0 \quad a_{\frac{n+3}{2}} \ \cdots \ a_n \\ R &= x \quad y \quad z \quad w \quad b_5 \ \cdots \ b_{\frac{n-1}{2}} \quad b_{\frac{n+1}{2}} \quad b_{\frac{n+3}{2}} \ \cdots \ b_n \\ S &= -x \quad -y \quad -z \quad -w \quad b_5 \ \cdots \ b_{\frac{n-1}{2}} \quad b_{\frac{n+1}{2}} \quad b_{\frac{n+3}{2}} \ \cdots \ b_n \end{aligned} \tag{3}$$

For the cases 1) to 5) use the values given in the following table or other suitable values:

$OD(4n; t, u, 4n - 18)$	t, u	p	q	r	s	x	y	z	w
	2	g	0	0	0	0	0	0	0
	4	g	0	0	0	g	0	0	0
	4, 4	g	h	0	0	g	$-h$	0	0
	2, 8	g	h	$-g$	0	g	0	g	0
	8, 8	g	g	h	$-h$	g	$-g$	h	h

(4)

3.2 The 5-2-2 Pattern

We also get the same theorem using the 5-2-2 pattern. Of course the results will be inequivalent.

Theorem 5 *Suppose there exists a $W(2n, 2n - 9)$ constructed using two circulants with first rows of the required pattern. Then there exist*

1. $OD(4n; 2, 4n - 18)$,
2. $OD(4n; 4, 4n - 18)$,
3. $OD(4n; 4, 4, 4n - 18)$,
4. $OD(4n; 2, 8, 4n - 18)$ and
5. $OD(4n; 8, 8, 4n - 18)$ constructed using four circulant matrices in the Goethals-Seidel array.

Proof We multiply each of the elements of the two $\{0, \pm 1\}$ sequences by the commuting variable a . We rearrange the required sequences of the $W(2n, 2n - 9)$, if necessary, to the form

$$\begin{aligned} A &= 0 \ 0 \ 0 \ 0 \ 0 \ a_6 \ \cdots \ a_n \\ B &= 0 \ 0 \ \star \ 0 \ 0 \ b_6 \ \cdots \ b_n \end{aligned} \tag{5}$$

where each element a_i and b_j is $\pm a$. The element \star is $\pm a$.

Now let p, q, r, s, x, y, z and w be commuting variables. Cases 1) to 5) are constructed by forming the four circulant matrices with first rows which are then used in the Goethals-Seidel array

$$\begin{aligned} P &= p \quad q \quad 0 \quad r \quad s \quad a_6 \quad \cdots \quad a_n \\ Q &= -p \quad -q \quad 0 \quad -r \quad -s \quad a_6 \quad \cdots \quad a_n \\ R &= x \quad y \quad \star \quad z \quad w \quad b_6 \quad \cdots \quad b_n \\ S &= -x \quad -y \quad \star \quad -z \quad -w \quad b_6 \quad \cdots \quad b_n \end{aligned} \tag{6}$$

For the cases 1) to 5) use the values given in the following table or other suitable values:

$OD(4n; t, u, 4n - 18)$	t, u	p	q	r	s	x	y	z	w
	2	g	0	0	0	0	0	0	0
	4	g	0	0	0	g	0	0	0
	4, 4	g	h	0	0	g	$-h$	0	0
	2, 8	g	h	$-g$	0	g	0	g	0
	8, 8	g	g	h	$-h$	g	$-g$	h	h

4 Inequivalent Hadamard matrices from the full $OD(4n; 2, 8, 4n - 10)$

The full orthogonal design $OD(4n; 2, 8, 4n - 10)$ constructed in theorem 3 can be used to construct Hadamard matrices of orders $4n$, for every n for which $W(2n, 2n - 5)$ weighing matrices are available. We carried out these computations for the values of n that belong to the set $\{15, 17, 21, 23, 25, 27, 29, 33, 35, 39, 43, 45, 47\}$. The corresponding $W(2n, 2n - 5)$ weighing matrices constructed from two circulants (many of

which are new) have been constructed in [4] and are also available in the web page <http://www.cargo.wlu.ca/weighing/>

For each value of n in $\{15, 17, 21, 23, 25, 27, 29, 33, 35, 39, 43, 45, 47\}$, we set the variables x, a, y to ± 1 values and for each of the resulting 8 combinations, we obtained a Hadamard matrix of order $4n$. Then we used the 4-profile criterion as implemented in Magma, to decide whether these Hadamard matrices are inequivalent. For each one of the thirteen orders $\{60, 68, 84, 92, 100, 108, 116, 132, 140, 156, 172, 180, 188\}$ we have located two inequivalent Hadamard matrices, having different 4-profiles. The results (the matrices and their 4-profiles) are available in the web page <http://www.cargo.wlu.ca/HMfromWM/>

In addition, one could check these Hadamard matrices for inequivalence, using the graph isomorphism criterion, which is more time consuming.

5 Conclusion

In this paper we have found new orthogonal designs using $W(2n, 2n - 5)$ and $W(2n, 2n - 9)$ weighing matrices constructed from two circulants. We also use the full $OD(4n; 2, 8, 4n - 10)$ to construct inequivalent Hadamard matrices of 13 orders.

References

- [1] R. Craigen, Weighing matrices and conference matrices, in *The CRC Handbook of Combinatorial Designs*, (Eds. C. J. Colbourn and J. H. Dinitz), CRC Press, Boca Raton, Fla., 1996, pp. 496–504.
- [2] R. Craigen and H. Kharaghani, Orthogonal designs, in *The CRC Handbook of Combinatorial Designs*, (Eds. C. J. Colbourn and J. H. Dinitz), 2nd ed., CRC Press, Boca Raton, Fla., 2006, pp. 290–306.
- [3] A. V. Geramita and J. Seberry, *Orthogonal designs. Quadratic forms and Hadamard matrices*, Lecture Notes in Pure and Applied Mathematics, 45, Marcel Dekker Inc. New York, 1979.
- [4] I. S. Kotsireas and C. Koukouvinos, New weighing matrices of order $2n$ and weight $2n - 5$, *J. Combin. Math. Combin. Comput.* (to appear).
- [5] I. S. Kotsireas, C. Koukouvinos and J. Seberry New weighing matrices of order $2n$ and weight $2n - 9$, submitted.
- [6] C. Koukouvinos and J. Seberry, New weighing matrices and orthogonal designs constructed using two sequences with zero autocorrelation function—a review, *J. Statist. Plann. Inference* 81 No. 1 (1999), 153–182.